

EFFICACY AND SAFETY OF TALETRECTINIB IN PATIENTS WITH ADVANCED OR METASTATIC ROS1+ NON-SMALL CELL LUNG CANCER: THE PHASE 2 TRUST-I STUDY


Wei Li,¹ Anwen Xiong¹, Nong Yang,² Huijie Fan,³ Qitao Yu,⁴ Yanqiu Zhao,⁵ Yongsheng Wang,⁶ Xue Meng,⁷ Jingxun Wu,⁸ Ziping Wang,⁹ Yunpeng Liu,¹⁰ Xicheng Wang,¹¹ Xintian Qin,¹² Kaihua Lu,¹³ Wu Zhuang,¹⁴ Yizhong Ren¹⁵, Xianyu Zhang¹⁵, Bing Yan,¹⁵ Christine M. Lovly,¹⁶ Caicun Zhou¹

¹Department of Medical Oncology, Shanghai Pulmonary Hospital and Thoracic Cancer Institute, Tongji University School of Medicine, Shanghai, China; ²Hunan Cancer Hospital, The Affiliated Cancer Hospital of Xiangya School of Medicine, Central South University, Changsha, China; ³The First Affiliated Hospital of Zhengzhou University, Zhengzhou, China; ⁴Medical Oncology of Respiratory, Affiliated Tumor Hospital of Guangxi Medical University, Nanning, China; ⁵Henan Cancer Hospital, Zhengzhou, China; ⁶West China Hospital Sichuan University, Chengdu, China; ⁷Department of Radiation Oncology, Shandong Cancer Hospital and Institute Jinan, China; ⁸The First Affiliated Hospital of Xiamen University, Xiamen, China; ⁹Beijing Cancer Hospital, Beijing, China; ¹⁰The First Hospital of China Medical University, Shenyang, China; ¹¹The First Affiliated Hospital/School of Clinical Medicine Guangdong Pharmaceutical University, Guangzhou, China; ¹²The First Affiliated Hospital of Guangdong Pharmaceutical University, Guangzhou, China; ¹³Jiangsu Province Hospital, Nanjing, China; ¹⁴Fujian Cancer Hospital, Fuzhou, China; ¹⁵Nuvation Bio, Inc., New York, NY, USA; ¹⁶Division of Hematology-Oncology, Vanderbilt University Medical Center and Vanderbilt Ingram Cancer Center, Nashville, TN, USA

Conflicts of Interest

I have no conflict of interest to declare.


TRUST-I (NCT04395677): Phase 2 Trial of Taletrectinib in ROS1+ NSCLC



- For patients with ROS1-positive (ROS1+) NSCLC, ROS1 tyrosine kinase inhibitors (TKIs) are the current standard of care¹
 - However, most patients progress within 2 years due to acquired resistance mutations (ie, G2032R) or brain metastases¹
- Taletrectinib is a next-generation, potent, CNS-active, and selective ROS1 TKI specifically designed to:^{2,3}
 - Improve efficacy and safety in ROS1+ NSCLC
 - Effectively treat CNS metastases
 - Overcome acquired resistance
 - Reduce neurological side effects with selectivity for ROS1 over tyrosine receptor kinase B (TrkB)
- We report updated data from the pivotal TRUST-I (NCT04395677) study in China, one of the largest clinical trials to date conducted in people living with ROS1+ NSCLC

1. Gendarme S et al. *Curr Oncol*. 2022;29:641-658. 2. Nagasaka M et al. *Future Oncol*. 2023;19(2):123-135. 3. Katayama R et al. *Nat Comm*. 2019;10(1):3604. CNS, central nervous system; NSCLC, non-small cell lung cancer; TKI, tyrosine kinase inhibitor.

TRUST-I (NCT04395677): Phase 2 Trial of Taletrectinib in ROS1+ NSCLC



Category	TKI Naive n=106	Crizotinib Pretreated n=67	Overall n=173
Median age, years (range)	56.0 (26–78)	51.0 (31–77)	55.0 (26–78)
Female, n (%)	59 (55.7)	41 (61.2)	100 (57.8)
ECOG PS 0/1, n (%)	20 (18.9)/86 (81.1)	19 (28.4)/48 (71.6)	39 (22.5)/134 (77.5)
Stage IV disease	97 (91.5)	65 (97.0)	162 (93.6)
Prior anticancer chemotherapy, n (%)	22 (20.8)	23 (34.3)	45 (26.0)
Never smoker	78 (73.6)	49 (73.1)	127 (73.4)
Brain metastasis, n (%)	18 (17.0)	28 (41.8)	46 (26.6)



^aSafety was analyzed in patients receiving ≥1 dose of taletrectinib until 30 days after the last dose of taletrectinib or the start date of new anticancer therapy minus 1 day, whichever occurred first.

^bA dose confirmation lead-in stage evaluated the safety of taletrectinib 400 mg QD (n=3) and 600 mg QD (n=3).

BOR, best overall response; cORR, confirmed objective response rate; DCR, disease control rate; DOR, duration of response; ECOG PS, Eastern Cooperative Oncology Group performance status; IC, intracranial; IRC, independent review committee; NSCLC, non-small cell lung cancer; PFS, progression-free survival; RECIST, Response Evaluation Criteria in Solid Tumors; TKI, tyrosine kinase inhibitor; TTR, time to response.

Taletrectinib: Efficacy in ROS1+ TKI-Naive NSCLC

Responses	TKI Naive n=106
IRC-assessed cORR, % (95% CI)	90.6 (83.33, 95.38)
DCR, % (95% CI)	95.3 (89.33, 98.45)
Median TTR, months (95% CI)	1.4 (1.38, 1.41)





Data cutoff: November 29, 2023. ^aTwo patients with confirmed BOR as not evaluable are not displayed in the waterfall and spider plots.

BOR, best overall response; cORR, confirmed objective response rate; CR, complete response; DCR, disease control rate; IRC, independent review committee; NSCLC, non-small cell lung cancer; PD, progressive disease; PR, partial response; SD, stable disease; TKI, tyrosine kinase inhibitor; TTR, time to response.

Taletrectinib: Efficacy in ROS1+ Crizotinib-Pretreated NSCLC

Responses	Crizotinib Pretreated n=66 ^a
IRC-assessed cORR, % (95% CI)	51.5 (38.88, 64.01)
DCR, % (95% CI)	83.3 (72.13, 91.38)
Median TTR, months (95% CI)	1.4 (1.38, 1.41)
cORR: G2032R mutations, % (n/N)	66.7 (8/12)





Data cutoff: November 29, 2023. ^aOne patient was excluded from the response-evaluable population in the crizotinib-pretreated group due to the presence of secondary cancer. ^bSix patients with confirmed BOR as not evaluable are not displayed in the waterfall and spider plots.
 BOR, best overall response; cORR, confirmed objective response rate; DCR, disease control rate; IRC, independent review committee; NSCLC, non-small cell lung cancer; PD, progressive disease; PR, partial response; SD, stable disease; TKI, tyrosine kinase inhibitor; TTR, time to response.

Taletrectinib: Duration of Response and Progression-Free Survival



TKI Naive (n=106)

Median follow-up: 23.5 mo



Crizotinib Pretreated (n=66)


Median follow-up: 9.7 mo



Data cutoff: November 29, 2023. DOR, duration of response; NR, not reached; PFS, progression-free survival; TKI, tyrosine kinase inhibitor.

Taletrectinib: Responses in Measurable Baseline Brain Metastases

Responses	TKI Naive n=8	Crizotinib Pretreated n=15
IC-cORR, % (95% CI)	87.5 (47.35, 99.68)	73.3 (44.90, 92.21)
DCR, % (95% CI)	100.0 (63.06, 100.0)	93.3 (68.05, 99.83)



Data cutoff: November 29, 2023. ^aOne patient with confirmed best overall response as not evaluable is not displayed in the waterfall plot.

cORR, confirmed objective response rate; DCR, disease control rate; IC, intracranial; IRC, independent review committee; PD, progressive disease; PR, partial response; SD, stable disease; TKI, tyrosine kinase inhibitor.

Taletrectinib Safety: TEAEs in ≥15% of Patients^a (N=173)

	Grade 1 n (%)	Grade 2 n (%)	Grade 3 n (%)	Grade 4 n (%)	Grade 5 ^b n (%)	Any grade n (%)
Increased AST	92 (53.2)	26 (15.0)	14 (8.1)	0	0	132 (76.3)
Diarrhea	99 (57.2)	16 (9.2)	6 (3.5)	0	0	121 (69.9)
Increased ALT	79 (45.7)	29 (16.8)	8 (4.6)	1 (0.6)	0	117 (67.6)
Vomiting	75 (43.4)	16 (9.2)	1 (0.6)	0	0	92 (53.2)
Anemia	52 (30.1)	30 (17.3)	3 (1.7)	0	0	85 (49.1)
Nausea	64 (37.0)	8 (4.6)	1 (0.6)	0	0	73 (42.2)
Decreased neutrophil count	25 (14.5)	10 (5.8)	6 (3.5)	4 (2.3)	0	45 (26.0)
Abnormal hepatic function	21 (12.1)	8 (4.6)	14 (8.1)	0	1 (0.6)	44 (25.4)
Decreased WBC count	27 (15.6)	14 (8.1)	3 (1.7)	0	0	44 (25.4)
Increased blood bilirubin	34 (19.7)	6 (3.5)	2 (1.2)	1 (0.6)	0	43 (24.9)
Dizziness	36 (20.8)	3 (1.7)	1 (0.6)	0	0	40 (23.1)
Proteinuria	34 (19.7)	5 (2.9)	0	0	0	39 (22.5)
Increased weight	17 (9.8)	16 (9.2)	3 (1.7)	0	0	36 (20.8)
Increased blood creatinine	33 (19.1)	2 (1.2)	0	0	0	35 (20.2)
QT prolongation	26 (15.0)	4 (2.3)	5 (2.9)	0	0	35 (20.2)
Hypercholesterolemia	29 (16.8)	4 (2.3)	0	0	0	33 (19.1)
Hyperuricemia	30 (17.3)	2 (1.2)	0	0	0	32 (18.5)
Decreased weight	23 (13.3)	8 (4.6)	0	0	0	31 (17.9)
Constipation	28 (16.2)	2 (1.2)	0	0	0	30 (17.3)
Decreased appetite	26 (15.0)	3 (1.7)	0	0	0	29 (16.8)
Increased conjugated bilirubin	22 (12.7)	3 (1.7)	2 (1.2)	1 (0.6)	0	28 (16.2)
COVID-19	10 (5.8)	15 (8.7)	3 (1.7)	0	0	28 (16.2)
Pyrexia	23 (13.3)	3 (1.7)	1 (0.6)	0	0	27 (15.6)
Increased blood CPK	21 (12.1)	5 (2.9)	0	0	0	26 (15.0)
Hypertriglyceridemia	24 (13.9)	2 (1.2)	0	0	0	26 (15.0)

- Median exposure of taletrectinib was 12.2 months (range: 0.23–40.04)
- 40.5% (70/173) of patients had a TEAE leading to treatment interruption
- 19.1% (33/173) of patients had a TEAE leading to a dose reduction
- 5.2% (9/173) of patients had a TEAE leading to treatment discontinuation
- Rates of neurologic TEAEs were low (dizziness: 23%; dysgeusia: 10%) and mostly grade 1

Data cutoff: November 29, 2023. ^aWorst grade per patient is reported. ^bTaletrectinib-related grade 5 TEAEs occurred in 3 patients: 2 TKI naive (1 hepatic failure, 1 pneumonia) and 1 crizotinib pretreated (abnormal hepatic function).

ALT, alanine aminotransferase; AST, aspartate aminotransferase; CPK, creatinine phosphokinase; TEAE, treatment-emergent adverse event; WBC, white blood cell.

Conclusions

- With long-term follow-up in the TRUST-I trial in China, taletrectinib continued to demonstrate meaningful efficacy in both TKI-naive and crizotinib-pretreated patients with ROS1+ NSCLC
 - High and durable responses were observed, including high intracranial ORR
 - Prolonged PFS was observed regardless of line of therapy
 - Taletrectinib demonstrated activity against acquired resistance mutations including G2032R
- Taletrectinib demonstrated a favorable safety profile
 - Rates of dose reduction and treatment discontinuation due to TEAEs continued to be low
 - The rate of neurologic TEAEs was low and most were grade 1
- The ongoing global pivotal phase 2 study, TRUST-II (NCT04919811), is further evaluating the efficacy and safety of taletrectinib in patients with ROS1+ NSCLC in the United States, Europe, and Asia¹

NSCLC, non-small cell lung cancer; ORR, objective response rate; PFS, progression-free survival; TEAE, treatment-emergent adverse event.

1. Nagasaka M et al. *Future Oncol*. 2023;19(2):123-135.

Efficacy and Safety of Taletrectinib in Chinese Patients With ROS1+ Non-Small Cell Lung Cancer: The Phase II TRUST-I Study



ascopubs.org



@JCO_ASCO



Acknowledgments

- We would like to thank all patients who participated in this study, the study investigators and their staff.
- This study was sponsored by AnHeart Therapeutics, a Nuvation Bio Company. Writing support was provided by David Jensen, PhD, of Peloton Advantage, LLC, an OPEN Health company, and was funded by Nuvation Bio.